Complete Summary

TITLE

Chronic stable coronary artery disease: percentage of patients queried one or more times about cigarette smoking.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement. Clinical performance measures. Chronic stable coronary artery disease. Chicago (IL): American Medical Association (AMA); 2003. 8 p.

Brief Abstract

DESCRIPTION

This measure assesses the percentage of patients with chronic stable coronary artery disease (CAD) who were queried one or more times about cigarette smoking.

RATIONALE

Smoking status should be determined and smoking cessation counseling and interventions are recommended (ACC/AHA/ACP-ASIM, 1999; ACC/AHA, 2000; ACC/AHA, 1999; ACC/AHA, 1999).

PRIMARY CLINICAL COMPONENT

Coronary artery disease; queried about cigarette smoking

DENOMINATOR DESCRIPTION

All patients with coronary artery disease (CAD)

NUMERATOR DESCRIPTION

The number of patients from the denominator who were queried one or more times about cigarette smoking

Evidence Supporting the Measure

PRIMARY MEASURE DOMAIN

Process

SECONDARY MEASURE DOMAIN

Not applicable

EVIDENCE SUPPORTING THE MEASURE

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

NATIONAL GUIDELINE CLEARINGHOUSE LINK

- 1999 update: ACC/AHA guidelines for the management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction).
- ACC/AHA guidelines for coronary artery bypass graft surgery: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1991 Guidelines for Coronary Artery Bypass Graft Surgery).
- ACC/AHA 2002 guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Unstable Angina).
- ACC/AHA 2002 guideline update for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for the Management of Patients With Chronic Stable Angina).

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Overall poor quality for the performance measured Wide variation in quality for the performance measured

EVIDENCE SUPPORTING NEED FOR THE MEASURE

Braunwald E, Antman EM, Beasley JW, Califf RM, Cheitlin MD, Hochman JS, Jones RH, Kereiakes D, Kupersmith J, Levin TN, Pepine CJ, Schaeffer JW, Smith EE 3d, Steward DE, Theroux P, Alpert JS, Eagle KA, Faxon DP, Fuster V, Gardner TJ, Gregoratos G, Russell RO, Smith SC Jr. ACC/AHA guidelines for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction. J Am Coll Cardiol 2000 Sep; 36(3):970-1062. [515 references] PubMed

Eagle KA, Guyton RA, Davidoff R, Ewy GA, Fonger J, Gott JP, Herrmann HC, Marlow RA, Nugent WC, O'Connor GT, Orsulak TA, Rieselbach RE, Winters WL,

Yusuf S, Gibbons RJ, Alpert JS, Garson A Jr, Gregoratos G, Russell RO, Smith SC Jr. ACC/AHA Guidelines for coronary artery bypass graft surgery. J Am Coll Cardiol 1999 Oct; 34(4): 1262-347.

Gibbons RJ, Chatterjee K, Daley J, Douglas JS, Fihn SD, Gardin JM, Grunwald MA, Levy D, Lytle BW, O'Rourke RA, Schafer WP, Williams SV, Ritchie JL, Cheitlin MD, Eagle KA, Gardner TJ, Garson A Jr, Russell RO, Ryan TJ, Smith SC Jr. ACC/AHA/ACP-ASIM guidelines for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 1999 Jun; 33(7): 2092-197. [891 references] PubMed

Jencks SF, Huff ED, Cuerdon T. Change in the quality of care delivered to Medicare beneficiaries, 1998-1999 to 2000-2001. JAMA 2003 Jan 15; 289(3): 305-12. PubMed

Ryan TJ, Antman EM, Brooks NH, Califf RM, Hillis LD, Hiratzka LF, Rapaport E, Riegel B, Russell RO, Smith EE III, Weaver WD. 1999 update: ACC/AHA guidelines for the management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 1999 Sep; 34(3):890-911. [849 references] PubMed

State of Use of the Measure

STATE OF USE

Pilot testing

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Patients of all ages with the diagnosis of chronic stable coronary artery disease.

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Not applicable

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

- Approximately 13 million Americans are living with coronary artery disease (CAD).
- More than 1 million Americans had a new or recurrent coronary attack in 2001.

EVIDENCE FOR INCIDENCE/PREVALENCE

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

ASSOCIATION WITH VULNERABLE POPULATIONS

Not applicable

BURDEN OF ILLNESS

- Chronic stable coronary artery disease (CAD) is the leading cause of mortality in the United States, accounting for almost 1 in 5 deaths.
- For individuals with CAD, the risk of another heart attack, stroke, and other serious complications is substantial.

EVIDENCE FOR BURDEN OF ILLNESS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

UTILIZATION

Within the past 2 decades, the number of short-stay hospital discharges for individuals with coronary artery disease (CAD) increased by almost 18%.

EVIDENCE FOR UTILIZATION

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

COSTS

The total annual cost of coronary artery disease (CAD) in the United States is approximately \$130 billion.

EVIDENCE FOR COSTS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

This performance measure is designed for prospective data collection in the office-based practice only. The measurement period may begin with the date of the most recent office visit, regardless of the diagnosis at that visit, and the data collection continues until 12 months are completed.

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR (INDEX) EVENT

Clinical Condition

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients with coronary artery disease (CAD)

Exclusions Unspecified

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients who were queried one or more times about cigarette smoking

Exclusions

None

DENOMINATOR TIME WINDOW

Time window follows index event

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

None

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Identifying Information

ORIGINAL TITLE

Smoking cessation.

MEASURE COLLECTION

The Physician Consortium for Performance Improvement Measurement Sets

MEASURE SET NAME

American College of Cardiology, American Heart Association, and Physician Consortium for Performance Improvement: Chronic Stable Coronary Artery Disease Core Physician Performance Measurement Set

SUBMITTER

American Medical Association on behalf of the American College of Cardiology, the American Heart Association, and the Physician Consortium for Performance Improvement

DEVELOPER

American College of Cardiology American Heart Association Physician Consortium for Performance Improvement

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2001 Aug

REVISION DATE

2003 Jan

MEASURE STATUS

This is the current release of the measure.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement. Clinical performance measures. Chronic stable coronary artery disease. Chicago (IL): American Medical Association (AMA); 2003. 8 p.

MEASURE AVAILABILITY

The individual measure, "Smoking Cessation," is published in the "Chronic Stable Coronary Artery Disease Core Physician Performance Measurement Set." This document is available from the American Medical Association (AMA) Division of Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.

For further information, please contact AMA staff by e-mail at cqi@ama-assn.org.

COMPANION DOCUMENTS

The following are available:

- Physician Consortium for Performance Improvement. Introduction to physician performance measurement sets. Tools developed by physicians for physicians. Chicago (IL): American Medical Association (AMA); 2001 Oct. 21 p. This document is available from the American Medical Association (AMA) Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.
- Physician Consortium for Performance Improvement. Principles for performance measurement in health care. A consensus statement. [online]. Chicago (IL): American Medical Association (AMA), Joint Commission on the Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); [3 p]. This document is available from the AMA Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.
- Physician Consortium for Performance Improvement. Desirable attributes of performance measures. A consensus statement. [online]. American Medical Association (AMA), Joint Commission on Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); 1999 Apr 19 [cited 2002 Jun 19]. [5 p]. This document is available from the AMA Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.

For further information, please contact AMA staff by e-mail at cgi@ama-assn.org.

NQMC STATUS

This NQMC summary was completed by ECRI on September 26, 2003.

COPYRIGHT STATEMENT

This Physician Performance Measurement Set (PPMS) was developed by the Physician Consortium for Performance Improvement (The Consortium) to facilitate quality improvement activities by physicians. The performance measures contained in this PPMS are not clinical guidelines and do not establish a standard of medical care. This PPMS is intended to assist physicians in enhancing quality of care and is not intended for comparing individual physicians to each other or for

individual physician accountability by comparing physician performance against the measure or guideline. The Consortium has not tested this PPMS.

This PPMS is subject to review and may be revised or rescinded at any time by The Consortium. The PPMS may not be altered without the prior written approval of The Consortium. A PPMS developed by The Consortium, while copyrighted, can be reproduced and distributed without modification for noncommercial purposes. Any other use is subject to the approval of The Consortium. Neither The Consortium nor its members shall be responsible for any use of this PPMS.

© 2003 American Medical Association

© 2004 National Quality Measures Clearinghouse

Date Modified: 11/1/2004



